arXiv:1701.07185v1 [math.RA] 25 Jan 2017

Nil extensions of Clifford ordered semigroup

A. K. Bhuniya and K. Hansda

Department of Mathematics, Visva-Bharati University,

Santiniketan-731235, West Bengal, India

anjankbhuniya@gmail.com, kalyanh4@gmail.com

January 26, 2017

Abstract

In this paper we describe all those ordered semigroups which are the nil extension of Clifford, left Clifford, group like, left group like ordered semigroups.

Key Words and phrases: nil extension, regular ordered semigroups, group like, left group like, Clifford and left Clifford ordered semigroups.

2010 Mathematics subject Classification: 20M10; 06F05.

1 Introduction:

Nil extensions of a semigroup (without order), are precisely the ideal extensions by a nil semigroup. In 1984, S. Bogdanović and S. Milić [2] characterized the semigroups (without order) which are nil extensions of completely simple semigroups, where as, a similar work was done by J. L Galbiati and M.L Veronesi [14] in 1980. Authors like S. Bogdanović, M. Cirić, Beograd have investigated this type extensions for regular semigroup, group, periodic semigroup as well as completely regular semigroup(see [4], [5]).

The notion of ideal extensions in ordered semigroups is actually introduced by N. Kehayopulu and M. Tsingelis [8]. In [11], they have worked on ordered semigroups which are nil extensions of Archimedean ordered semigroups. The concepts of nil extensions have been extended to ordered semigroups by Y. Cao [7], with characterizing ordered semigroups which are nil extensions of simple,

left simple, t-simple ordered semigroups. Further he described complete semilattices of nil extensions of such ordered semigroups.

The aim of this paper is to describe all those ordered semigroups which are the nil extension of Clifford, left Clifford, group like, left group like ordered semigroups. This research originates from the research papers [4], [5].

Our paper organized as follows. The basic definitions and properties of ordered semigroups are presented in Section 2. Section 3 is devoted to characterizing the nil extensions of Clifford and left Clifford ordered semigroups.

2 Preliminaries:

In this paper $\mathbb{N}$ will provide the set of all natural numbers. An ordered semigroup is a partiality ordered set S, and at the same time a semigroup $(S, \cdot)$ such that (forall $a,\ b,\ x \in S$) $a \leq b \Rightarrow xa \leq xb$ and $ax \leq bx$. It is denoted by $(S, \cdot, \leq)$. For an ordered semigroup S and $H \subseteq S$, denote $(H] := \{t \in S : t \leq h,\ \text{for some } h \in H\}$. An element z in S is called zero of S if $z < x$ and $zx = xz = z$ for every $x \in S$.

Let I be a nonempty subset of an ordered semigroup S. I is a left (right) ideal of S, if $SI \subseteq I$ ($IS \subseteq I$) and $(I] = I$. I is an ideal of S if I is both a left and a right ideal of S. An (left, right) ideal $I\ of S$ is proper if $I \neq S$. The intersection of all ideals of an ordered semigroup S, if nonempty, is called the kernel of S and is denoted by $K(S)$.

An ordered semigroup S is called a group like ordered semigroup if for all $a, b \in S$ there are $x, y \in S$ such that $a \leq xb$ and $a \leq by$ [1] and S left group like if for all $a, b \in S$ there is $x \in S$ such that $a \leq xb$ [1]. Kehayopulu [9] defined Greens relations on an ordered semigroup S as follows: $a\mathcal{L}b\ if\ L(a) = L(b),\ a\mathcal{R}b$ if $R(a) = R(b),\ a\mathcal{J}b$ if $I(a) = I(b)$, and $\mathcal{H} = \mathcal{L} \cap \mathcal{R}$. These four relations $\mathcal{L},\ \mathcal{R},\ \mathcal{J}$ and $\mathcal{H}$ are equivalence relations. In an ordered semigroup S, an equivalence relation ρ is called left (right) congruence if for $a, b, c \in S\ a\ \rho\ b$ implies $ca\ \rho\ cb$ ($ac\ \rho\ bc$). ρ is congruence if it is both left and right congruence. A congruence ρ on S is called semilattice if for all $a, b \in S\ a\ \rho\ a^2$ and $ab\ \rho\ ba$. A semilattice congruence ρ on S is called complete if $a \leq b$ implies $(a, ab) \in \rho$. The ordered semigroup S is called complete semilattice of subsemigroup of type τ if there exists a complete semilattice congruence ρ such that $(x)_\rho$ is a type τ subsemigroup of S. Equivalently: There exists a semilattice Y and a family of subsemigroups $\{S\}_{\alpha \in Y}$ of type τ of S such that:

1. $S_\alpha \cap S_\beta = \phi$ for any $\alpha,\ \beta \in Y$ with $\alpha \neq \beta$,

2. $S = \bigcup_{\alpha \in Y} S_\alpha$,
3. $S_\alpha S_\beta \subseteq S_{\alpha\,\beta}$ for any $\alpha,\ \beta \in Y$,
4. $S_\beta \cap (S_\alpha] \neq \phi$ implies $\beta \preceq \alpha$, where $\preceq$ is the order of the semilattice Y defined by $\preceq := \{(\alpha,\ \beta) \mid \alpha = \alpha\ \beta\ (\beta\ \alpha)\}$ [10].

Let S be an ordered semigroup with the zero 0. An element $a \in S$ is called a nilpotent if $a^n = 0$ for some $n \in \mathbb{N}$. The set of all nilpotents of S is denoted by $Nil(S)$. S is called nil ordered semigroup (nilpotent) [7] if $S = Nil(S)$.

Due to Cao [7] the definition of nil extension of ordered semigroup is as follows.

Definition 2.1. *[7] Let I be an ideal of an ordered semigroup S. Then $(S/I, \cdot, \preceq)$ is called the Rees factor ordered semigroup of S modulo I, and S is called an ideal extension of I by ordered semigroup S/I. An ideal extension S of I is called a nil-extension of I if S/I is a nil ordered semigroup.*

Lemma 2.2. *[7] Let S be an ordered semigroup and I an ideal of S. Then the following are equivalent:*

(i) S is a nil-extension of I;

(ii) $(\forall a \in S)(\exists m \in \mathbb{N})\ a^m \in I$.

In [1], we have introduced the notion of Clifford and left Clifford ordered semigroups and characterized their structural representation. For the correspondences of the results of this paper we are stating some of them.

Definition 2.3. *[1] A regular ordered semigroup S is called Clifford ordered (left Clifford ordered) semigroup if for all $a, b \in S$ and $ab \in (bSa]$ $(ab \in (Sa])$.*

Theorem 2.4. *[1] Let S be a regular ordered semigroup. Then followings hold in S:*

1. *S is Clifford if and only if $\mathcal{L} = \mathcal{R}$.*
2. *$\mathcal{L}$ is a complete semilattice congruence if S is Clifford.*
3. *S is Clifford ordered semigroup if and only if it is a complete semilattice of group like ordered semigroups.*

Theorem 2.5. *[1] Let S be a regular ordered semigroup. Then followings hold in S:*

1. *$\mathcal{L}$ is a complete semilattice congruence if S is left Clifford.*
2. *S is left Clifford ordered semigroup if and only if it is a complete semilattice of left group like ordered semigroups.*

3 Main Results:

Now we describe all those ordered semigroups which are the nil extensions of Clifford, left Clifford, group like, left group like ordered semigroups. We omit the proof of the following lemma as it is straightforward.

Lemma 3.1. *An ordered semigroup S is left group like ordered semigroup if and only if $a \in (aSab]$ if for all $a,\ b \in S$.*

Theorem 3.2. *An ordered semigroup S is a nil extension of a left group like ordered semigroup if and only if for every $a, b \in S$, there exists $n \in \mathbb{N}$ such that $a^n \in (a^nSa^nb]$ and for every $a \in S,\ b \in Reg_{\leq}(S),\ a \leq ba$ implies $a \leq axb$ for some $x \in S$.*

Proof. Suppose that S is a nil extension of a left group like ordered semigroup K and $a, b \in S$. Then there is $m \in \mathbb{N}$ such that $a^m \in K$. Regularity of K implies that $a^m \leq a^mxa^m$ for some $x \in K$. Further, for $xa^m, a^mb \in K$; the left simplicity of K yields that $xa^m \leq ya^mb$, for some $y \in K$. This gives that $a^m \leq a^mxa^m \leq a^mya^mb$. Next let $b \in Reg_{\leq}(S)$ and $a \in S$ such that $a \leq ba$. Since $b \in Reg_{\leq}(S)$, there exists $z \in S$ such that $b \leq b(zb)^n$ for all $n \in \mathbb{N}$. Then for some $n_1 \in \mathbb{N}$, $(zb)^{n_1} \in K$; whence $b(zb)^{n_1} \in K$. Thus $b \in K$ and so $ba \in K$. Thus $a \in K$. Since K is a left group like ordered semigroup, for $a, ba \in K$ it follows that $a \leq asba$ for some $s \in K$, by Lemma 3.1. Thus the given conditions follow.

Conversely, assume that given conditions hold in S. Let $a \in S$ be arbitrary. Then by given condition we have $a^m \leq a^mxa^{m+1}$, for some $x \in S$ and $m \in \mathbb{N}$. This implies $a^{m+1} \in Reg_{\leq}(S)$ and so $Reg_{\leq}(S) \neq \phi$. Denote $T = Reg_{\leq}(S)$.

Let us choose $s \in S$ and $a \in T$. Then the definition of T implies

$$a \leq a(xa)^n, \text{ for all } n \in \mathbb{N} \text{ and some } \ x \in S.$$

Thus $sa \leq sa(xa)^n$, for all $n \in \mathbb{N}$. Now for $xa, sa \in S$, there exists $m_1 \in \mathbb{N}$ and $t_1 \in S$ such that

$$(xa)^{m_1} \leq (xa)^{m_1}t_1(xa)^{m_1}sa, \text{ by the first condition.}$$

Then $a \leq a(xa)^{m_1}$ implies $sa \leq sa(xa)^{m_1}$ and hence $sa \leq sa(xa)^{m_1}t_1(xa)^{m_1}sa$; where $(xa)^{m_1}t_1(xa)^{m_1} \in S$. So $sa \in T$. Also $a \leq (ax)^na$ for all $n \in \mathbb{N}$. Let $m_2 \in \mathbb{N}$ be such that $(ax)^{m_2} \in T$. Then $as \leq (ax)^{m_2}as$ implies by the second condition that

$$as \leq ast_2(ax)^{m_2} \text{ for some } t_2 \in S. \tag{1}$$

Denote $t = (ax)^{m_2}$. Then the definition of T implies

$$t \leq t(zt)^n \text{ for all } n \in \mathbb{N} \text{ for some } \ z \in T. \tag{2}$$

Then using the first condition for $zt, as \in S$ we have that $(zt)^{m_3} \leq (zt)^{m_3} t_3 (zt)^{m_3} as$ for some $m_3 \in \mathbb{N}$ and $t_3 \in S$. That is $t \leq t(zt)^{m_3} t_3 (zt)^{m_3} as$, by (2). So from (1) we have $as \leq ast_2 t(zt)^{m_3} t_3 (zt)^{m_3} as$, and hence $as \in T$.

Next choose $a \in S$ and $b \in T$ such that $a \leq b$. Since $b \in T$ there is $x \in S$ such that $b \leq b(xb)^n$ for all $n \in \mathbb{N}$. Now for $xb, a \in S$, it follows from the first condition that

$$(xb)^{m_4} \leq (xb)^{m_4} t_4 (xb)^{m_4} a \text{ for some } \ m_4 \in \mathbb{N} \text{ and } t_4 \in S. \tag{3}$$

So $a \leq b$ implies that

$$\begin{aligned} a &\leq b(xb)^{m_4} t_4 (xb)^{m_4} a \\ &= bt_5 a; \text{ where } t_5 = (xb)^{m_4} t_4 (xb)^{m_4}. \end{aligned}$$

Since $b \in T$, by above we have $\ bt_5 \in T$. Say $bt_5 = t_6$. Then for $a \in S$, and $t_6 \in T$, $\ a \leq t_6 a$ yields that $a \leq at_7 t_6$ for some $t_7 \in S$, by second condition. Therefore

$$\begin{aligned} a &\leq at_7 bt_5 \\ &= at_7 b(xb)^{m_4} t_4 (xb)^{m_4} \\ &\leq at_7 b(xb)^{m_4} t_4 (xb)^{m_4} t_4 (xb)^{m_4} a, \text{ by (3)}. \end{aligned}$$

Clearly $t_7 b(xb)^{m_4} t_4 (xb)^{m_4} t_4 (xb)^{m_4} \in T$, as $b \in T$. Thus $a \in T$, which shows that T is an ideal of S.

Finally let $c, d \in T$. Then there is $v \in S$ such that $c \leq c(vc)^n$ for all $n \in \mathbb{N}$. Now for $vc, d \in S$, there exists $t_8 \in S$ such that $c \leq c(vc)^{m_5} t_8 (vc)^{m_5} d$ for some $m_5 \in \mathbb{N}$. Since $c \in T$, $c(vc)^{m_5} t_8 (vc)^{m_5} \in T$. Hence T is left simple. Thus T is left group like ordered semigroup such that for every $a \in S$ there is $m \in \mathbb{N}$, $a^m \in T$. Hence S is nil extension of a left group like ordered semigroup T. □

In the following result we provide an independent proof of Corollary 5.2 of [7].

Theorem 3.3. *An ordered semigroup S is a nil extension of a group like ordered semigroup if and only if for all $a, b \in S$, there exists $n \in \mathbb{N}$ such that $a^n \in (b^n S b^n]$.*

Proof. Suppose that S is a nil extension of a group like ordered semigroup G and $a, b \in S$. Then there exists $n \in \mathbb{N}$ such that $a^n, b^n \in G$. Since G is a group like ordered semigroup, there exists $u \in G$ such that $a^n \leq b^n u$. Also for $u, b^n \in G$ there exists $x \in G$ such that $u \leq xb^n$. This implies $b^n u \leq b^n x b^n$. Thus $a^n \leq b^n x b^n$ and hence $a^n \in (b^n S b^n]$.

Conversely, let us assume that given condition holds in S. Choose $a \in S$. Then for some $m \in \mathbb{N}$ and $x \in S$, $a^m \leq a^m x a^m$. Thus $Reg_{\leq}(S) \neq \phi$. Say $G = Reg_{\leq}(S)$. So for every $a \in S$, there exists $m \in \mathbb{N}$ such that $a^m \in G$. Let us consider $b \in G$ and $s \in S$. Then for all $n \in \mathbb{N}$

$$bs \leq bybs \leq (by)^n bs \text{ for } y \in S. \tag{4}$$

Using the given condition for $bs, by \in S$, we obtain $(by)^m bs \leq (bs)^m z (bs)^{m+1}$ for $z \in S$ and $m \in \mathbb{N}$. This yields that

$$\begin{aligned} bs &\leq (bs)^m z (bs)^{m+1} \\ &\leq bstbs; \text{ where } t = bs^{m-1} z b s^m. \end{aligned}$$

Thus $bs \in G$. Similarly $sb \in G$.

Next let $a \in S$ and $b \in G$ be such that $a \leq b$. Since $b \in G$ there exists $x \in S$ such that $b \leq bxb$ and hence $b \leq (bx)^n b (xb)^n$ for all $n \in \mathbb{N}$, which implies that

$$\begin{aligned} a &\leq a^m (z_1 a^m b a^n z) a^n, \text{ for some } m, n \in \mathbb{N} \text{ and } z, z_1 \in S \\ &= ata, \text{ where } t = a^{m-1} z_1 a^m b a^n z a^{n-1}. \end{aligned}$$

So $a \in G$. Hence G is an ideal of S.

Finally, consider $a, b \in G$. Then there exists $x \in S$ such that

$$a \leq (ax)^n a \text{ for all } n \in \mathbb{N},$$

and so by the given condition it follows that $a \leq b^{m'} z' b^{m'} a$ for some $m' \in \mathbb{N}$ and $z' \in S$. This gives that $a \leq bu$ for some $u = b^{m'-1} z' b^{m'} a \in G$. Similarly there is some $v \in S$ such that $a \leq vb$. This shows that G is a group like ordered semigroup. Hence S is a nil extension of a group like ordered semigroup G. □

Theorem 3.4. *An ordered semigroup S is a nil extension of a Clifford ordered semigroup if and only if for every $x, a, y \in S$, there exists $n \in \mathbb{N}$ such that $xa^n y \in (xa^n ySya^n x] \cap (ya^n xSxa^n y]$ and $a \in S$, $b \in Reg_{\leq}(S)$ such that $a \leq b$, implies $a \in (Sab]$.*

Proof. First suppose that S is a nil extension of a Clifford ordered semigroup K. Let $x, a, y \in S$. Then there is $m \in \mathbb{N}$ such that $a^m \in K$. Since K is an ideal of S, $xa^m y \in K$. Since K is a regular, there exists $z_1 \in K$ such that

$$xa^m y \leq xa^m y z_1 x a^m y. \tag{5}$$

Now $z_1 x, a^m y \in K$ implies that

$$z_1 x a^m y \leq (a^m y) u_1 (z_1 x), \text{ for some } u_1 \in K, \text{ since } S \text{ Clifford.} \tag{6}$$

Similarly for $a^m, (y u_1 z_1) \in K$ there is $u_2 \in S$ such that

$$a^m (y u_1 z_1) \leq (y u_1 z_1) u_2 a^m. \tag{7}$$

Therefore

$$\begin{aligned} xa^m y &\leq xa^m y z_1 x a^m y \\ &\leq xa^m y a^m (y u_1 z_1) x, \text{ by (6)} \\ &\leq xa^m y^2 u_1 z_1 u_2 a^m x, \text{ by (7).} \end{aligned}$$

Thus

$$xa^m y \leq xa^m y z_1 x a^m y^2 u_1 z_1 u_2 a^m x. \tag{8}$$

Also, for $a^m y^2, u_1 z_1 u_2 \in K$ there exists $u_3 \in K$ such that $a^m y^2 u_1 z_1 u_2 \leq u_1 z_1 u_2 u_3 a^m y^2$. Then from (8), we obtain that

$$\begin{aligned} xa^m y &\leq xa^m y (z_1 x u_1 z_1 u_2 u_3 a^m y) y a^m x \\ &\leq xa^m y s y a^m x; \text{ where } s = z_1 x u_1 z_1 u_2 u_3 a^m y. \end{aligned}$$

Therefore $xa^m y \in (xa^m y S y a^m x]$. Similarly $xa^m y \in (ya^m x S x a^m y]$.

Now $K \subseteq Reg_{\leq}(S)$ implies that $Reg_{\leq}(S) \neq \phi$. Consider $b \in Reg_{\leq}(S)$ and $a \in S$ such that $a \leq b$. Since $b \in Reg_{\leq}(S)$, there exists $z \in S$ such that $b \leq (bz)^n b$ for all $n \in \mathbb{N}$. Since S is a nil extension of K, there exists $n_1 \in \mathbb{N}$ such that $(bz)^{n_1} \in K$. This gives $(bz)^{n_1} b \in K$, which gives $b \in K$ and so $a \in K$, since K is an ideal of S. Since K is a Clifford ordered semigroup, by Theorem 2.4(ii) $\mathcal{L}$ is a congruence on S. Since $a, b \in K$ we have $a \mathcal{L} ab$ and hence $a \subseteq (Sab]$.

Conversely, let us assume that given conditions hold in S. Let $a \in S$ be arbitrary. Then by the first condition there exists $n \in \mathbb{N}$ such that $a^{n+2} \leq a^{n+2} x a^{n+2}$,

for some $x \in S$. Thus $Reg_{\leq}(S) \neq \phi$. Say $T = Reg_{\leq}(S)$. It is now clear that for each $a \in S$, there exists $m \in \mathbb{N}$ such that $a^m \in T$.

Let $s \in S$ and $x \in T$. Then for all $n \in \mathbb{N}$ and for some $t \in S$, $x \leq (xt)^n x$ which implies that $sx \leq sx(tx)^{n-1}tx$, for all $n \in \mathbb{N}$. By first condition there are $s_1 \in S$ and $m_1 \in \mathbb{N}$ such that $sx \leq sx(tx)^{m_1}txs_1tx(tx)^{m_1}sx$ and thus $sx \leq sxpsx$; where $p = (tx)^{m_1}txs_1tx(tx)^{m_1}$. Also for every $n \in \mathbb{N}$,

$$\begin{aligned} xs &\leq x(tx)^n s \\ &\leq xt(xt)^{n-1}xs. \end{aligned}$$

So there is $m_2 \in \mathbb{N}$ such that

$$\begin{aligned} xs &\leq xs(xt)^{m_2}xts_2xt(xt)^{m_2}xs \\ &\leq xsqxs; \text{ where } q = (xt)^{m_2}xts_2xt(xt)^{m_2}. \end{aligned}$$

Thus $sx, sx \in T$.

To show T, a Clifford ordered semigroup, choose $a, b \in T$. Then there is $r \in S$ such that

$$\begin{aligned} ab &\leq abrab \\ &\leq (abra)(bra)^{n-1}b, \text{ for all } n \in \mathbb{N}. \end{aligned} \tag{9}$$

Now for $abra, bra, b \in S$, the first condition yields that

$$abra(bra)^{m_3}b \leq b(bra)^{m_3}abrap_1abra(bra)^{m_3}b \text{ for some } p_1 \in S \text{ and } m_3 \in \mathbb{N}.$$

Therefore from (9)

$$\begin{aligned} ab &\leq b(bra)^{m_3}abrap_1abra(bra)^{m_3}b \\ &\leq b(bra)^{m_3}abrap_1abra(bra)^{m_3-1}brab \\ &\leq bgab; \text{ where } g = (bra)^{m_3}abrap_1abra(bra)^{m_3-1}br \in T. \end{aligned} \tag{10}$$

Similarly there are $m_4 \in \mathbb{N}$ and $p_2 \in S$ such that $ab \leq abra(bra)^{m_4}bp_2b(bra)^{m_4}abra$. So from (10),

$$ab \leq b(gabra(bra)^{m_4}bp_2b(bra)^{m_4})abra = bua; \text{ where } u = gabra(bra)^{m_4}bp_2b(bra)^{m_4}abr \in T.$$

Hence T is Clifford ordered semigroup.

Now let $a \leq b$ for some $a \in S$ and $b \in T$. Then by the second condition, there is $z \in S$ such that $a \leq zab$, that is,

$$a \leq zabtab \text{ for some } t \in S. \tag{11}$$

Since T is Clifford ordered semigroup, for $zabt, ab \in T$ it follows that

$$zabtab \leq abp_3zabt \text{ for some } p_3 \in S.$$

Similarly for $bp_3za, bt \in T$, we have $bp_3zabt \leq btp_4bp_3za$ for some $p_4 \in S$.

The last two inequalities together with (11) yields that $a \leq aha$, where $h = btp_4bp_3z$. Thus $a \in T$ and so T is an ideal of S. Hence S is a nil extension of a Clifford ordered semigroup T. □

Theorem 3.5. *An ordered semigroup S is a nil extension of a left Clifford ordered semigroup if and only if for every $x, a, y \in S$, there exists $n \in \mathbb{N}$ such that $xa^ny \in (xa^nySya^nx] \cap (xa^nySxa^ny]$ and $a \in S$, $b \in Reg_{\leq}(S)$ such that $a \leq b$ implies $a \leq azab$ for some $z \in S$.*

Proof. Let S be a nil extension of a left Clifford ordered semigroup K. Choose $x, a, y \in S$. Then there exists $m \in \mathbb{N}$ such that $a^m \in K$. Since K is an ideal of S, $xa^my \in K$. Also the regularity of K yields that

$$xa^my \leq xa^myz_1xa^my, \text{ for some} z_1 \in K. \tag{12}$$

Since K is a left Clifford ordered semigroup and $z_1x, a^my \in K$, by Theorem 2.5 it follows that $z_1xa^my \leq z_2(z_1x)$ for some $z_2 \in K$. Therefore

$$\begin{aligned} xa^my &\leq xa^myz_1xa^my \\ &\leq xa^myz_1xa^myz_1xa^my \\ &\leq xa^myz_1xa^myz_2(z_1x). \end{aligned} \tag{13}$$

Similarly, for $a^m, yz_2z_1 \in K$ there is $z_3 \in K$ such that

$$a^myz_2z_1 \leq z_3a^m, \tag{14}$$

and for $a^m, z_1xz_3 \in K$ there is $z_4 \in K$ such that

$$a^myz_1xz_3 \leq z_4a^my. \tag{15}$$

Thus from (13) we obtain that

$$\begin{aligned} xa^my &\leq xa^myz_1xa^myz_2z_1x \\ &\leq xa^myz_1xa^mz_3a^mx, \text{ from (14)} \\ &\leq xa^myz_4a^mya^mx, \text{ from (15)}. \end{aligned}$$

Hence $xa^my \in (xa^mySya^mx]$ and so $xa^my \in (xa^mySya^mx] \cap (xa^mySxa^my]$, from (12).

To show the second condition choose $a \in S$ and $b \in Reg_{\leq}(S)$ be such that $a \leq b$. By the regularity of b yields that $b \leq b(tb)^n$ for some $t \in S$ and for all $n \in \mathbb{N}$. Then there is $r \in \mathbb{N}$ such that $(tb)^r \in K$. Since K ia an ideal $b(tb)^r \in K$ and so $b \in K$. Thus $a \in K$. Since K is left Clifford ordered semigroup, $\mathcal{L}$ is congruence on S, by Theorem 2.5(i). Thus $a\mathcal{L}ab$. Then a, ab are in Theorem 3.1, $a \leq azab$ for some $z \in S$. This proves the necessary condition.

Conversely, suppose that given conditions hold in S. Let $a \in S$. Then by the first condition there exists $n \in \mathbb{N}$ such that $a^{n+2} \leq a^{n+2}xa^{n+2}$, for some $x \in S$. Thus $Reg_{\leq}(S) \neq \phi$. Say $T = Reg_{\leq}(S)$. Now for each $a \in S$, there exists $m \in \mathbb{N}$ such that $a^m \in T$. Let $s \in S$ and $x \in T$. Then for all $n \in \mathbb{N}$ and for some $t \in S$, $x \leq (xt)^n x$. This implies $sx \leq sx(tx)^{n-1}tx$, for all $n \in \mathbb{N}$.

Then by the first condition there are $s_1 \in S$ and $m_1 \in \mathbb{N}$ such that

$$\begin{aligned} sx &\leq sx(tx)^{m_1}txs_1tx(tx)^{m_1}sx \\ &\leq sxpsx; \text{ where } p = (tx)^{m_1}txs_1tx(tx)^{m_1}. \end{aligned}$$

Therefore $sx \in T$.

We now show that T is a left Clifford ordered semigroup. For this let us assume that $a, b \in T$. Then there is $t_1 \in S$ such that

$$ab \leq abt_1ab \leq a(bt_1a)^nbt_1ab \text{ for all } n \in \mathbb{N}.$$

Then by first condition there are $t_2 \in S$ and $m' \in \mathbb{N}$ such that

$$\begin{aligned} ab &\leq a(bt_1a)^{m'}bt_1ab \\ &\leq a(bt_1a)^{m'}bt_1abt_2bt_1ab(bt_1a)^{m'}a \end{aligned}$$

Therefore $ab \leq t_1'a$; where $t_1' = a(bt_1a)^{m'}bt_1abt_2bt_1ab(bt_1a)^{m'}$. Since $a \in T$ we have $t_1' \in T$. Hence T is left Clifford ordered semigroup.

Next let $a \in S$ and $b \in T$ such that $a \leq b$. Using second condition we have $a \leq azab$, for some $z \in S$. Then for all $n \in \mathbb{N}$, $a \leq a(za)^nb^n$. So for some $m'' \in \mathbb{N}$, $(za)^{m''} \in T$. Since T is a left Clifford, $(za)^{m''}b^{m''} \in (Ta]$. So $a \in (aSa]$, that is $a \in T$.

Finally to show T, an ideal of S we need only to show that $xs \in T$. The regularity of x yields that $xs \leq x(tx)^ns$ for all $n \in \mathbb{N}$. Also for some $l \in \mathbb{N}$, $x(tx)^ls \in T$. Then $xs \leq x(tx)^ls$ implies that $xs \in T$ by above. Thus T is an ideal of S. Hence S is a nil extension of a left Clifford ordered semigroup T. □

References

[1] A.K.Bhuniya and K. Hansda, Complete semilattice of ordered semigroups, Communicated.

[2] S. Bogdanović and S. Milić, A nil extension of completely simple semigroup, *Publications de L'Institut, Nouvelle Serie*, **36(50)** (1984).

[3] S. Bogdanović, Semzgroups with a system of subsemigroups, *Inst. of Math , Novi Sad*, 1985.

[4] S. Bogdanović and M. Cirić, A nil extension of regular semigroup, *Glasnik Matmaticki*, **25(2)** (1991).

[5] S. Bogdanovć, M. Cirić, Semilattices of Archimedean semigroups and (completely) pi-regular semigroups, I (A survey), *Filomat (Ni)* **7** (1993), 1-40.

[6] S. Bogdanovićand M. Cirić, Semilattices of nil extension of simple regular semigroup, *Algebra Colloquium*,**10(1)**(2003), 81-90.

[7] Y. Cao and X. Xinzhai, Nil extensions of simple po-semigroups, *Communication in Algebra*, **28(5)**(2000), 2477-2496.

[8] N. Kehayopulu and M. Tsingelis, Ideal extensions of ordered semigroups, *Comm. Algebra*, **31(10)**(2003), 4939-4969.

[9] N.Kehayopulu, Ideals and Green's relations in ordered semigroups, *International Journal of Mathematics and Mathematical Sciences* , (2006), 1-8, Article ID 61286.

[10] N. Kehayopulu and M. Tsingelis, Semilattices of Archimedean Ordered Semigroups, *Algera Colloquium* **15:3**(2008), 527-540.

[11] N.Kehayopulu, Archimeadean ordered semigroups as ideal extensions,*Semigroup Forum* , **78**(2009), 343-348.

[12] M.Petrich, Lectures in Semigroups, *John Wiley and Sons* , 1977.

[13] D. Kees, On semigroups, *Proc. Cambridge Phil. Soc.*, **36** (1940), 387-400.

[14] J. L Galbiati and M.L Veronesi, Sui semiguppi che sono un band di t-semigruppi, *Instituto Lombardo (Rend. Sc)*, **A 114** (1980), 217-234.